

Gauss-Bonnet type theorems in any dimension ^{*}

Valery DOLOTIN[†]

Institute of Theoretical & Experimental Physics, Moscow

February 1, 2008

Abstract

Given a construction of smooth homotopy class invariants of smooth immersions $M^n \rightarrow \mathbf{R}^{n+k}$. The particular case of $k = 1$, $n \geq 1$ is a sequence of non-zero integrals, where the $n = 2$ term is the Gauss-Bonnet integral

1 Basic example

Take a smooth curve $\gamma = (x(t), y(t)) \subset \mathbf{R}^2$.

Statement 1.1

$$I(\gamma) := \int \frac{\dot{y}\ddot{x} - \dot{x}\ddot{y}}{\dot{x}^2 + \dot{y}^2} dt = 2\pi$$

Proof Denote $u := \dot{x}$, $v := \dot{y}$. Then the integral becomes

$$\int_{\gamma} \frac{vdu - u dv}{u^2 + v^2}$$

where $\frac{vdu - u dv}{u^2 + v^2}$ is a closed form on $\mathbf{R}^2 - 0$ being a generator of $H^1(\mathbf{R}^2 - 0; \mathbf{R})$.
□

Note, that the above statement is the smooth analogue of the Euclidean geometry theorem about the sum of exterior angles of a polygon.

If the curve has self-intersections, then in general $I(\gamma) = 2\pi k$ where k is the "winding number" of the tangent vector while moving along the curve. If we take a homotopy connecting two curves with different winding numbers then we have to pass through a curve with a singularity of the form $\frac{vdu - u dv}{u^2 + v^2}$ at some point. Those singular curves make a discriminant set in the space of smooth curves.

2 General construction

2.1 Codimension 1 case

For each space $P = (p_0, \dots, p_n)$ we have a canonical closed form

$$\omega := \frac{\sum_i (-1)^i p_i dp_0 \wedge \dots \wedge \widehat{dp_i} \wedge \dots \wedge dp_n}{(p_0^2 + \dots + p_n^2)^{(n+1)/2}}$$

^{*}This work was partially supported by RFBR 01-01-00546 grant.

[†]e-mail: vd@gate.itep.ru

on $P - 0$.

Then for a smooth immersion of n -manifold $M \rightarrow \mathbf{R}^{n+1}$ in each local chart $(t_1, \dots, t_n)$ on M we have a Jacobian $n \times (n+1)$ matrix

$$J := \left(\frac{\partial x_i}{\partial t_j} \right)$$

If we substitute via Plucker embedding $p_i = \Delta_i := \Delta_{0 \dots \hat{i} \dots n}(J)$ (being the minor build on all except i -th columns of J) into ω then we get an integration form $\omega(\Delta)$ on M . This form is correctly defined since the orientation preserving change of coordinates between charts multiplies all $\Delta_{\hat{i}}$ by the same number (the Jacobian of the change) and ω is invariant under such multiplication. For a non-orientable manifold the form $\omega(\Delta)$ may be interpreted as a two-valued form, or a form on a double covering of M .

Since ω is closed then

Statement 2.1

$$\int_M \omega(\Delta) = 2^n \pi k$$

where k is an integer invariant of a smooth homotopy class of smooth immersions of M .

Note that in codimension 1 case this invariant equals to the degree of the corresponding Gauss map $M^n \rightarrow S^n$.

2.1.1 Example: Gauss-Bonnet theorem

Here $n = 2$ and

$$\omega = \frac{p_0 dp_1 \wedge dp_2 - p_1 dp_0 \wedge dp_2 + p_2 dp_0 \wedge dp_1}{(p_0^2 + p_1^2 + p_2^2)^{3/2}}$$

For a smooth immersion $M^2 \rightarrow \mathbf{R}^3$, $(t_1, t_2) \mapsto (x_1, x_2, x_3)$ with the Jacobian

$$J = \left(\frac{\partial x_i}{\partial t_j} \right) =: (x_{i,j})$$

and $p_0 = \Delta_0(J) = x_{1,1}x_{2,2} - x_{2,1}x_{1,2}$, $p_1 = \Delta_1(J) = x_{0,1}x_{2,2} - x_{2,1}x_{0,2}$, $p_2 = \Delta_2(J) = x_{0,1}x_{1,2} - x_{1,1}x_{0,2}$ we have the integral

$$\int_M \omega(\Delta) = \int_M \frac{\sum_{i,j} \Delta_i \Delta_j dx_{i,1} \wedge dx_{j,2}}{(\sum \Delta_i^2)^{3/2}} = \int \frac{\sum_{i,j} \Delta_i \Delta_j (x_{i,11}x_{j,22} - x_{i,12}x_{j,21}) dt_1 dt_2}{(\sum \Delta_i^2)^{3/2}}$$

One can check that $\sum \Delta_i^2 = EG - F^2$ is the determinant (in Gauss notations) of the metric form induced on M from the Euclidean metric on $\mathbf{R}^3$. Now for the numerator

$$\begin{aligned} & \sum_{i,j} \Delta_i \Delta_j (x_{i,11}x_{j,22} - x_{i,12}x_{j,21}) = \\ & = \begin{vmatrix} x_{1,11} & x_{2,11} & x_{3,11} \\ x_{1,1} & x_{2,1} & x_{3,1} \\ x_{1,2} & x_{2,2} & x_{3,2} \end{vmatrix} \begin{vmatrix} x_{1,22} & x_{2,22} & x_{3,22} \\ x_{1,1} & x_{2,1} & x_{3,1} \\ x_{1,2} & x_{2,2} & x_{3,2} \end{vmatrix} \end{aligned}$$

$$- \begin{vmatrix} x_{1,12} & x_{2,12} & x_{3,12} \\ x_{1,1} & x_{2,1} & x_{3,1} \\ x_{1,2} & x_{2,2} & x_{3,2} \end{vmatrix} \begin{vmatrix} x_{1,21} & x_{2,21} & x_{3,21} \\ x_{1,1} & x_{2,1} & x_{3,1} \\ x_{1,2} & x_{2,2} & x_{3,2} \end{vmatrix} = ln - m^2$$

is the determinant of the second fundamental form (in Gauss notations). Since the area element on M is $dS = \sqrt{EG - F^2} dt_1 dt_2$ then we have

$$\int_M \omega(\Delta) = \int \frac{ln - m^2}{(EG - F^2)^2} dS = 2\pi\chi(M)$$

which is the Gauss-Bonnet formula.

2.2 Affine Gauss map

For a smooth immersion $\phi : M^n \rightarrow \mathbf{R}^{n+k}$ taking the minors of its Jacobian we get a map $\mathcal{J} : M^n \rightarrow G_{n,n+k}$, which is the higher codimension analogue of the Gauss map.

Then for any closed n -form $\omega \in H^n(G_{n,n+k})$ its pullback $\mathcal{J}^*\omega$ gives us a smooth homotopy class invariant $\int_M \mathcal{J}^*\omega$ of smooth immersions of M .

The generators of $H^n(G_{n,n+k})$ have a natural representation in terms of Plucker coordinates p_I on $G_{n,n+k}$ as

$$\omega = \frac{\sum \varphi_{I_1 \dots I_n}(p) dp_{I_1} \wedge \dots \wedge dp_{I_n}}{|p|^d}$$

where $|p| := (\sum p_I^2)^{1/2}$, and $\varphi_{I_1 \dots I_n}(p)$ are homogeneous polynomials of degree $d - n$.

2.2.1 Example: $n = 2$, Kahler form

On the space $\mathbf{R}^{2N} - \{0\}$ we have a closed 2-form

$$\begin{aligned} \omega = & \left(\sum_{i,j} (p_i dp_j + p_{i+N} dp_{j+N}) \right) \wedge (p_{j+N} dp_j - p_j dp_{j+N}) \\ & - (p_j dp_i + p_{j+N} dp_{i+N}) \wedge (p_{i+N} dp_i - p_i dp_{i+N}) / |p|^4 \end{aligned}$$

which is the imaginary (non-zero) part of the Kahler form $\partial\bar{\partial} \log(|z_1|^2 + \dots + |z_N|^2)$ written in terms of complex coordinates $z_j = p_j + ip_{j+N}$.

Note, that as a form on $\mathbf{R}^{2N} - \{0\}$ it is exact, but it is invariant under the natural $\mathbf{C}^*$ -action on $\mathbf{R}^{2N} - \{0\}$ and its pull-back to the space $\mathbf{C}P^{N-1}$ of $\mathbf{C}^*$ -orbits gives a non-trivial cohomology class.

In particular, for a smooth complex submanifold $X^n \subset \mathbf{C}^{n+k}$ the corresponding Jacobian matrix, combined with the Plucker embedding gives us a map $X^n \rightarrow \mathbf{C}P^{\binom{n}{n+k}-1}$. Then the pull-back of ω is known as the first Chern class $c_1(X \cap H)$ of hyperplane section line bundle over X . On the other hand, taking X^n as a real $2n$ -dimensional immersion into $\mathbf{R}^{2(n+k)}$ we have the pull-back of ω in terms of real partial derivatives $\partial x_i / \partial t_j$, $i = 1, \dots, 2(n+k)$, $j = 1, \dots, 2n$. Since the real d -closeness of ω is just the algebraic property of its expression in terms of these real partial derivatives, then $\int_M J^* \omega^n$ is a differential invariant for any immersion $M^{2n} \rightarrow \mathbf{R}^{2(n+k)}$.

2.3 Projective Gauss map

For each point x of a smooth projective manifold $M^n \subset \mathbf{R}P^N$ we have a $(n+1)$ -dimensional subspace in $\mathbf{R}^{N+1}$, tangent to M at x , given by the Jacobian matrix of the cone over M in $\mathbf{R}^{N+1}$. So we have a (projective) Gauss map $J : M^n \rightarrow G_{n+1, N+1}$.

Example 2.1 Take $n = 1$, $N = 2$, i.e. a curve $\gamma \subset \mathbf{R}P^2$. Take a chart $U_0 := \{(x_0, x_1, x_2) \mid x_0 \neq 0\} \subset \mathbf{R}^3 - \{0\}$ and write the 2-parametric equation of the cone over γ as $x_0 = s$, $x_1 = sf_1(t)$, $x_2 = sf_2(t)$. The corresponding Jacobian matrix is

$$J = \begin{pmatrix} 0 & sf'_1 & sf'_2 \\ 1 & f_1 & f_2 \end{pmatrix}$$

Take the form $\omega_0 = \frac{p_2 dp_1 - p_1 dp_2}{p_1^2 + p_2^2}$ on $\mathbf{R}^3 - \{0\}$. Then its pull-back $J^*\omega_0$ to the image of Plucker embedding $p_i = \Delta_i(J)$ of the Gauss image of γ is the form $\frac{f_2 \dot{f}_1 - f_1 \dot{f}_2}{f_1^2 + f_2^2} dt$, known from Section 1.

Note, that if γ has a singularity in the U_0 chart then $J^*\omega_0$ has a singularity at the corresponding point of γ (although the inverse is not true). Then we have a bundle $\omega(\alpha) := \sum_0^2 \alpha_i \omega_i$ of singular forms on $\mathbf{R}^3$ of which the pull-backs to γ give integrals, being invariants if smooth homotopy classes of the Gauss map $\gamma \rightarrow \mathbf{R}P^2 - \{q_0, q_1, q_2\}$, where $q_i = \{p \in \mathbf{R}P^2 \mid \sum_{j \neq i} p_j^2 = 0\}$.

In the chart $U_i := \{(x_0, \dots, x_N) \mid x_i \neq 0\}$ the cone over M may be given parametrically as $x_0 = s \cdot f_0(t_1, \dots, t_n), \dots, x_i = s, \dots, x_N = s \cdot f_N(t_1, \dots, t_n)$ with the Jacobian

$$J = \begin{pmatrix} sf_{0,t_1} & \dots & 0 & \dots & sf_{N,t_1} \\ \vdots & & \vdots & & \vdots \\ sf_{0,t_n} & \dots & 0 & \dots & sf_{N,t_n} \\ f_0 & \dots & \underbrace{1}_i & \dots & f_N \end{pmatrix}$$

where $f_{i,t_j} := \frac{\partial f_i}{\partial t_j}$. Having a singularity of M in the chart U_i implies setting $\Delta_{ii_1 \dots i_n}(J) = 0$, for all $i_1 < \dots < i_n \subset \{1, \dots, N\}$, which also implies the degeneracy of the whole J . Then on the space of Plucker coordinates $\mathbf{R}^{\binom{n+1}{N+1}}$ any form

$$\omega = \left(\sum \varphi_{I_1 \dots I_n}(p) dp_{I_1} \wedge \dots \wedge dp_{I_n} \right) / |p|^d$$

(where $\varphi(p)$ are homogeneous of degree $d - n$) has a pull-back $J^*\omega$ to M with singularities at all singular points of M , regardless the chart of their location.

Now take ω to be closed.

Statement 2.2 For the pull-back $J^*\omega$ to M the integral

$$\int_M J^*\omega$$

is a smooth homotopy class invariant of $M^n \subset \mathbf{R}P^N$.

Note, that even if ω is exact as a form on $\mathbf{R}^{\binom{n+1}{n+1}} - \{0\}$ its pull-back to the projective manifold may give a non-trivial class (as it's the case in the above example).

For a given immersion $M^n \rightarrow \mathbf{R}^{n+k}$ we may take a composition of the affine Gauss map $\mathcal{J}_a : M^n \rightarrow G_{k,n+k} \hookrightarrow \mathbf{R}P^{\binom{k}{n+k}-1}$ with the projective one $\mathcal{J}_p : \mathcal{J}_a(M) \rightarrow G_{n+1, \binom{k}{n+k}}$ and use the corresponding invariants.

3 Gauss-Bonnet type invariants for polyhedra

$n = 1$ This is the above mentioned theorem on the sum of the exterior angles of a polygon.

$n = 2$ For a vertex ν of a triangulation the sum of angles over all triangles τ of its star

$$\sum_{\tau \in St_\nu} (\alpha_\tau + \beta_\tau + \gamma_\tau) = \pi |St_\nu|$$

(where $|St_\nu|$ is the number of triangles) is a constant, invariant with respect to different immersions of the star. If α_τ denote angles adjacent to ν then we can rewrite this identity as

$$2\pi - \sum_{\tau} \alpha_\tau = 2\pi - \sum_{\tau} (\pi - (\beta_\tau + \gamma_{\tau+1}))$$

Note, that $\pi - (\beta_\tau + \gamma_{\tau+1})$ may be interpreted as a deformed exterior angle at a vertex $\mu \in Ln_\nu$ with adjacent β_τ and $\gamma_{\tau+1}$ which differs from the actual exterior angle as "much" as the immersion of the star at μ differs from lying in a 2-plane. We call this difference by a curvature $K_1(\mu)$ of a 1-dimensional link at the vertex μ . Since 2π is the topological invariant of the link, which is equal to the sum of exterior angles when the link lies in a 2-plane then we can rewrite the above equation as

$$E_2(\nu) := 2\pi - \sum_{\tau \in St_\nu} \alpha_\tau = 2\pi - \sum_{\mu \in Ln_\nu} K_1(\mu)$$

and call $E_2(\nu)$ the **exterior 2-angle** of ν via the immersion into $\mathbb{R}^3$. Now the analogue of the theorem on exterior angles of a polygon is

Statement 3.1

$$\sum_{\nu} E_2(\nu) = 2^2 \pi \kappa$$

is the topological invariant of the immersion.

Inductive step

Having defined the exterior angle $E_k(\nu)$ for a vertex of a k -polyhedron in $\mathbb{R}^{k+1}$ and knowing the corresponding topological invariant $2^k \pi$ of a k -sphere we can define the curvature $K_k(\mu)$ at vertices of the k -link of a $k+1$ -star immersed into $\mathbb{R}^{k+2}$. Then the $k+1$ -exterior angle is defined by induction

$$E_{k+1}(\nu) := 2^k \pi - \sum_{\mu \in Ln_\nu} K_k(\mu)$$

which may be regarded as a particular case of polyhedral Stokes formula, $2^k\pi - E_{k+1}(\nu)$ corresponding to integration form of degree $k+1$.

The corresponding invariant of immersions into $\mathbb{R}^{k+2}$ is

$$\sum_{\nu \in P} E_{k+1}(\nu)$$

4 Discussion

The above consideration implies, that having a differential invariant of a manifold, written in terms of "intrinsic geometry" of M (such as Gauss-Bonnet), it may be convenient to represent it as a restriction onto M of some natural closed form defined on the appropriate target space of a smooth immersion of M .

It would be interesting to see the polyhedral interpretation of the integral $\int_M \omega(\Delta)$ for $\text{codim } M \geq 2$ analogous to that in the curvature case above. Provided having the polyhedral presentation of $\omega(\Delta)$ we could compute the value of the corresponding polyhedral form at vertices of any polyhedra, including non-smoothable ones. Since each polyhedron may be smoothed almost everywhere then each embedding of such almost everywhere smooth manifold has irremovable singularities at certain points corresponding to "non-smoothable" vertices of our triangulation. Then the value of the polyhedral form at non-smoothable vertices should distinguish them among others, giving some maximal possible value (like 2π value for the curvature form at the cusp singularity of a plane curve).